

Certificate of Analysis

COA No: CA_XBE-0021-2

Version: 04

Lyo-Compatible MMLV-RT

Suitable for Research and further Manufacturing Use

Catalog No:	MDX042
Lot No:	EN055-B345360
Storage Conditions:	-20°C
Component Lot No.	LCR-225107B
Expiry date:	August 2027

Quality Control Parameters

High-concentration MMLV-RT suitable for incorporation into lyophilized RT-PCR assays

Analysis	Specification	Result
Functional	Activity is measured as reverse transcriptase units by quantitative PCR analysis against a reference enzyme. Pass Criteria: Activity must be greater than 165 U/μL	966.0 U/μL
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in concordance with control sample. Pass Criteria: Amplification traces must overlay with the negative control.	Passed
DNase contamination	DNase contamination is measured as DNA substrate degradation against a DNase I dilution series by agarose gel electrophoresis. Limit of detection: 6.25 x 10 ⁻⁴ KU DNase I. Pass Criteria: No detectable degradation.	Passed



Certificate of Analysis

COA No: CA_XBE-0021-2

Version: 04

RNase contamination	Quantitative PCR analysis with high and low RNase standards. Limit of detection: 9.7 x 10 ⁻³ ng/µL RNase Pass Criteria: Test sample must show less RNase activity than the limit of detection.	Passed
---------------------	--	--------

QA / QC Representative:

J. Rahnenführer

Date: 15th July 2025

United Kingdom

Tel: +44 (0)20 8830 5300 Fax: +44 (0)20 8452 2822 USA

Germany

Tel: +49 (0)3371 60222 00 Tel: +1 901.382.8716 Fax: +1 901.382.0027 Fax: +49 (0)3371 60222 01



Certificate of Analysis

COA No: CA_BDB-0025-2

Version: v05

Enzyme Dilution Buffer

For research or further manufacturing use only

Catalog No:	MDX042	
Lot No:	EN055-B345360	
Storage Conditions:	-20°C	
Lot number:	TDB-525207B	
Expiry date:	August 2027	

Quality Control Parameters

Enzyme Dilution Buffer is a glycerol-free 10x buffer used for the dilution of Taq antibody or reverse transcriptase.

Analysis	Specification	Result
	A 3Kb fragment was amplified with a dilution series of Enzyme Dilution Buffer, using standard conditions and 30 cycles.	
Functional	A 3Kb fragment was amplified with a dilution series of <i>Taq polymerase</i> , using standard conditions and 30 cycles.	Passed
	Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).	
Glycerol determination	Glycerol content is < 0.2% determined by spectrophotometric analysis and comparison to a standard curve.	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 6.25 x 10^{-4} KU/ μ L.	Passed
RNase contamination	Quantitative PCR analysis with high and low RNase standards. Test sample must show less RNase activity than the limit of detection: 9.7x10 ⁻³ ng/µl RNase.	Passed

QA / QC Representative:

J. Rahnenführer

Date: 15th July 2025

United Kingdom

Tel: +44 (0)20 8830 5300 Fax: +44 (0)20 8452 2822 USA

Germany

Tel: +1 901.382.8716 Tel: +49 (0)3371 60222 00 Fax: +49 (0)3371 60222 01 Fax: +1 901.382.0027